INDEX

Acaroletis pseudococci, 148–49.
Adalia bipunctata, 348, 441.
Ædes cantator, 260.
  sollicitans, 260.
  sylvestris, 264, 265.
Ægerita webberi, 475.
Ageniaspis fuscicollis, 297.
Agonoderma pallipes, 477.
Agrilus bilineatus, 373.
  fagi, 438.
  politus, 438–40.
  sinuatus, 251.
  viridis, 438.
Agriotes mancus, 308.
Agromyza pusilla, 447–54.
  scutellata, 447–54.
Æolothrips fasciatus, 217.
Ainslie, G. G., 322–24.
Alabama argillacea, 152–53, 405.
Alcohol, commercial, 272.
Aldrich, J. M., 404.
Allograpta fracta, 295.
  obliqua, 295.
Ammonia as a fumigant, 149–50.
Amphorophora rubicola, 136.
Anaphothrips striatus, 211–18.
Angumois grain moth, 204.
Anti-mosquito work, 260–68.
Apanteles lateralis, 187.
Aphanogmus floridanus, 330.
Aphelinus diaspidis, 227.
  fuscipennis, 227.
Aphidius testaceipes, 295.
Aphis atriplicis, 133.
  bakeri, 133.
  brassicæ, 440, 441, 443.
  frigidæ, 132.
  medicaginis, 295, 296.
  viburni, 440, 441, 443.
Apiary inspection, 92–97.
Apple tent caterpillar, 373.
Applied entomology, expenditures, 57; growth, 50–64; scope, 60; today's work, 64.
Argentine ant, 147, 153.
Arsenate of lead, properties of various forms, 162–67.
Arthrocnodax carolina, 330.
  constricta, 481.
  rhoina, 481.
Aschersonia aleyrodis, 475.
  flavocitrina, 475.
Aspidiotus camelliæ, 468.
  cyanophylli, 468.
  cydoniæ, 468.
  hederæ, 468.
  juglans-regiæ, 468.
  perniciosus, 422.
  tangæ, 250–51.
Atmometers, 249.
Aulacaspis pentagona, 298.
  rosæ, 468.

Baccha clavata, 295, 296.
Barbey, A., 350.
Bartonia bacilliformis, 357.
Bee-keeping in Missouri, 96–97.
Bee moth, observations on, 183–88.
Bichromate of potash, 269, 273.
Black leaf, 40, 195.
  weevil, 204.
Blissus leucopterus, 414, 415.
Boll weevil, 65–67, 234–40.
Botrytis bassiana, 474.
Bourne, A. I., 196–200.
Bracon brevicornis, 187.
Britton, W. E., 244–46, 257–60.
Brown-tail moth, 250.
Brown-tail and gipsy moth work, 83–87, 116–119.
Burgess, A. F., 7, 83.
Busck, August, 340–41.

Cabbage aphis, 440.
Cadelle beetles, 318.
Calandra oryza, 204, 206–10, 315.
Canadian entomological service, 298–99.
  nursery regulations, 151.
Cane beetle borer, 390–98.
Canis caraibicus, 359, 360, 362, 363, 364.
  criollus, 362.
Carbolic acid emulsion, 199.
  and lime, 199.
Carbon bisulfide, 188, 198.
Carpocapsa pomonella, 177.
Cathartus advena, 316.
  gemellatus, 204, 317.
Cavia cabaya, 361.
Cebus capuchinus, 358.
Cecidomyia viticola, 339.
Cereal products, 313.
Ceresa bubalus, 241.
Ceromasia sphenophori, 390, 454.
Ceroplastes sp., 469.
Chaitophorus sp., 128.
Chelonus texanus, 296, 297.
Chestnut borer, two-lined, 373.
Chiggers, 152.
Chinch bug, 414, 415, 416.
  parasite, 219–22.
Chittenden, F. H., 152.
Chloride of lime, 285, 286, 287.
Chrysomphalus sp., 468.
Chrysopa oculata, 330.
Clean culture, 332.
Clerid larva, 251.
Clover leaf weevil, 297.
Cobalt, 273.
Coccidæ of Peru, 466.

INDEX

Acaroletis pseudococci, 148–49.
Adalia bipunctata, 348, 441.
Ædes cantator, 260.
  sollicitans, 260.
  sylvestris, 264, 265.
Ægerita webberi, 475.
Ageniaspis fuscicollis, 297.
Agonoderma pallipes, 477.
Agrilus bilineatus, 373.
  fagi, 438.
  politus, 438–40.
  sinuatus, 251.
  viridis, 438.
Agriotes mancus, 308.
Agromyza pusilla, 447–54.
  scutellata, 447–54.
Æolothrips fasciatus, 217.
Ainslie, G. G., 322–24.
Alabama argillacea, 152–53, 405.
Alcohol, commercial, 272.
Aldrich, J. M., 404.
Allograpta fracta, 295.
  obliqua, 295.
Ammonia as a fumigant, 149–50.
Amphorophora rubicola, 136.
Anaphothrips striatus, 211–18.
Angumois grain moth, 204.
Anti-mosquito work, 260–68.
Apanteles lateralis, 187.
Aphanogmus floridanus, 330.
Aphelinus diaspidis, 227.
  fuscipennis, 227.
Aphidius testaceipes, 295.
Aphis atriplicis, 133.
  bakeri, 133.
  brassicæ, 440, 441, 443.
  frigidæ, 132.
  medicaginis, 295, 296.
  viburni, 440, 441, 443.
Apiary inspection, 92–97.
Apple tent caterpillar, 373.
Applied entomology, expenditures, 57; growth, 50–64; scope, 60; today's work, 64.
Argentine ant, 147, 153.
Arsenate of lead, properties of various forms, 162–67.
Arthrocnodax carolina, 330.
  constricta, 481.
  rhoina, 481.
Aschersonia aleyrodis, 475.
  flavocitrina, 475.
Aspidiotus camelliæ, 468.
  cyanophylli, 468.
  cydoniæ, 468.
  hederæ, 468.
  juglans-regiæ, 468.
  perniciosus, 422.
  tangæ, 250–51.
Atmometers, 249.
Aulacaspis pentagona, 298.
  rosæ, 468.

Baccha clavata, 295, 296.
Barbey, A., 350.
Bartonia bacilliformis, 357.
Bee-keeping in Missouri, 96–97.
Bee moth, observations on, 183–88.
Bichromate of potash, 269, 273.
Black leaf, 40, 195.
  weevil, 204.
Blissus leucopterus, 414, 415.
Boll weevil, 65–67, 234–40.
Botrytis bassiana, 474.
Bourne, A. I., 196–200.
Bracon brevicornis, 187.
Britton, W. E., 244–46, 257–60.
Brown-tail moth, 250.
Brown-tail and gipsy moth work, 83–87, 116–119.
Burgess, A. F., 7, 83.
Busck, August, 340–41.

Cabbage aphis, 440.
Cadelle beetles, 318.
Calandra oryza, 204, 206–10, 315.
Canadian entomological service, 298–99.
  nursery regulations, 151.
Cane beetle borer, 390–98.
Canis caraibicus, 359, 360, 362, 363, 364.
  criollus, 362.
Carbolic acid emulsion, 199.
  and lime, 199.
Carbon bisulfide, 188, 198.
Carpocapsa pomonella, 177.
Cathartus advena, 316.
  gemellatus, 204, 317.
Cavia cabaya, 361.
Cebus capuchinus, 358.
Cecidomyia viticola, 339.
Cereal products, 313.
Ceresa bubalus, 241.
Ceromasia sphenophori, 390, 454.
Ceroplastes sp., 469.
Chaitophorus sp., 128.
Chelonus texanus, 296, 297.
Chestnut borer, two-lined, 373.
Chiggers, 152.
Chinch bug, 414, 415, 416.
  parasite, 219–22.
Chittenden, F. H., 152.
Chloride of lime, 285, 286, 287.
Chrysomphalus sp., 468.
Chrysopa oculata, 330.
Clean culture, 332.
Clerid larva, 251.
Clover leaf weevil, 297.
Cobalt, 273.
Coccidæ of Peru, 466.

Coccinella 9-notata, 442, 443.
transversoguttata, 441, 443.
trifasciata, 441, 442, 443.
Coccinellid statistics, 440–43.
Coccus hesperidum, 469.
Cockerell, T. D. A., 148, 405, 459.
Codling moth, 251.
Colorado potato beetle, 152.
Conical grape gall, 339.
Confused flour beetle, 318.
Conotrachelus nenuphar 168.
Cooley, R. A., 193–95.
Coquillettidia perturbans, 264.
Cordyceps militaris, 474.
Corn (stored), reducing insect injury, 203–10.
bill bug, southern, 83.
root worm, western, 322–24.
Correspondence filing, 87–91.
Cotton, 327.
worm moth, 405.
Crosby, C. R., 408.
Culex pipiens, 257–60.
salinarius, 265.
Currant fruit fly, 193.
Currants, two new pests, 193–95.
fruit weevil, 193–94.
Current notes, 158, 254, 301, 354, 409, 484.
Cycloneda sanguinea, 441, 442, 443.

Davidson, W. M., 127–136.
Dean, G. A., 67–73.
Derostenus dístatæ, 453.
Dew, J. A., 162–67.
Diabrotica longicornis, 477.
12 punctata, 477.
Diapheromera femorata, 299.
Diaspis boisduvalii, 469.
echinocacti, 470.
Diplazon lætatorius, 294–97.
sycophanta, 295.
Dissosteira carolina, 82.
Drug store beetle, 317.

Eccoptogaster quadrispinosa, 374–75.
Eciton schmitti, 147.
Economic entomology, bibliography, 34–36.
Editorial, 154, 253, 300, 349, 407, 482.
Elis sexcincta, 388.
Empria fragariæ, 480.
Empusa sphærosperma, 297.
Entomological investigations, 7–22.
publications, efficiency, 26–34.
work in Missouri, 376–78.
Entomologists association, constitution amended, 38.
education of, 274–80.
employment bureau, 5.
executive committee report, 4.
incorporation, 22–24.
list of meetings, ix; members, xii.
membership committee, 44–45.
nominations, 46.
Entomologists association, secretary's report, 2.
26th annual meeting, proceedings, 1–97, 162–233, 257–94, 305–39, 357–78.
Entomophthora aulicæ, 475.
Ephestia kuehniella, 319.
Epicauta marginata, 477.
vittata, 477.
Epochra canadensis, 193.
Escherich, K., 350–52.
Eucallipterus arundicolens, 129.
Euceraphis betulæ, 129.
Eumicrosoma benefica, 220.
Eupachylomma rileyi, 403–404.
Eupelmus cereanus, 187.
Eupeodes volucris, 295.
Euproctis chrysorrhœa, 177.
Euthrips nervosus, 212.
striatus, see Anaphothrips.
Evaporation, studies of, 229–33.
Evetria buoliana, 340–41.
Ewing, H. E., 440–43.
Exoprosopa fascipennis, 387.

Felt, E. P., 148–49, 173, 339, 373–75, 458–59, 481.
Fernald, H. T., 196–200, 228–29.
Fiorinia fioriniæ, 470.
Flint, W. P., 476.
Flour beetle, confused, 204, rust-red, 204.
Forest tent caterpillar, 373.
insects, 373–75.
Formalin, 272.
Forestiere, Traite d'Entomologie, 350.
Forstinsekten Mitteleuropas, 350–52.
Fink, D. E., 149–50, 400–401.
Fruit insects, manual, 408.
Fungous diseases, 473–76.

Galleria mellonella, 183–88.
Gates, B. N., 348.
Gibson, Arthur, 201–203.
Gipsy and brown-tail moth work, 83–87, 116–19.
Gipsy moth, 250, 368–73; larvæ, 170; parasite search, incident, 378–82.
Girault, A. A., 445–47.
Glaser, R. W., 473–76.
Goodwin, W. H., 313–22.
Gooseberries, two new pests, 193–95.
Gossard, H. A., 31.
Grain beetle, sawtoothed, 204.
square-necked, 204,
louse, southern, 414, 415.
Grass thrips, 211–18.
Grasshopper baits, 68, 76, 78–81.
control, 67–73.
Green fruit worm, 189–91.
soldier bug, 336–39.

Hammar, A. G., 155.
Haseman, L., 96–97, 124–27, 376–78.
Headlee, T. J., 260–68, 413–17.

Heliothis obsoleta, 175, 405, 477.
Heilipus lauri, 243–44.
Hellebore, 198.
Hemerocampa definita, 370.
rulosa, 370.
leucostigma, 252, 370.
Hemichionaspis minor, 469.
Hemlock borer, spotted, 373.
Herrick, G. W., 189–92, 483.
Hewitt, C. G., 151, 195, 211–18, 281–89.
Hickory bark beetle, 373–75.
Hinds, W. E., 65–67, 203–10.
Hippodamia convergens, 441, 442, 443.
spuria, 441, 442, 443.
Hodgkiss, H. E., 227–28.
Hop aphis, 440.
Household insects, 483.
Horticultural inspectors, proceedings, 98–127, 234–44; certificates, 342–48.
House fly, 281–89, 268–73, 289–93.
Howard, L. O., 247–48, 274–80, 378–82.
Hunter, S. J., 293–94.
and Claassen, 73–82.
Hunter, W. D., 234–40, 289–92.
Hypera punctata, 297.
Hypoderma bovis, 195.
Hyslop, J. A., 305–12.

Icerya zeteki, 148.
Illingworth, J. F., 390–98, 444–45.
Incorporation, Entomologists, 22–24.
Indian meal moth, 204, 318.
Insect metabolism and moisture, 413–17.
resistant to sprays, 167–72.
Insectary, experience with, 181–83.
Inspection (nursery) administration, 119–23.
certificates standards, 342–48.
laws, 102–106.
notes, 240–44.
work, 124–27.
plant diseases, 106–108.
Iridomyrmex humilis, 147, 153.
Iron sulphate, 285, 287.
Isosoma grande, 403.
Itonididæ, zoöphagous, 458–59.

Jones, T. H., 460–62.

Kaliosysphinga dohrnii, 251.
Karschomyia cocci, 461, 462.
Kelly, E. O. G., 294–97.
Kermes waldeni, 150–51.
Kerosene emulsion, 199, 286, 287.
Killifish, 262.
King, J. L., 401–403.
King, G. B., 150–51, 246–47.

Lachnosterna, 382, 386, 389.
Læmophlæus minutus, 317.
Laphygma frugiperda, 461.
Lasius niger americanus, 476–78.
Leaf hoppers in Maine, economic notes, 192.
Lepidosaphes beckii, 471.
Leptinotarsa decemlineata, 152, 177.
Lepus cuniculus, 359.
Leucopis flavicornis, 404.
Lime-sulphur wash, 167–71, 195.
valuation, 463–67.
Limnerium disparidis, 378.
Limothrips avenæ, 217.
Liothrips montanus, 194–95.
Lithomyza condita, 460.
Lysiphlebus tritici, 414, 415.
Lissorhoptrus simplex, 433–39.
Litomastax truncatellus, 297.
Loxostege similalis, 296.

Macrosiphum ludovicianæ, 136.
Malacosoma americana, 175, 373.
disstria, 373.
Matheson, R., 141–47.
Mason, P. W., 252.
McColloch, J. W., 219–27.
McGregor, E. A., 324–36, 447–54.
Microweisia misella, 431.
Megilla maculata, 348.
Melanoplus atlanis, 67.
bivittatus, 67, 82.
differentialis, 67, 82.
Melander, A. L., 167–72.
Melanophila fulvoguttata, 373.
Mephitis mephitis putida, 252.
Meromyza americana, 212.
Merrill, D. E., 251–52.
Mesogramma polita, 295, 296.
Metcalf, C. L., 218.
Metcalf, Z. P., 83.
Microcerata iridis, 460.
Milk and formalin, 274.
Milk, sweet, 273.
Miscible oils, 173.
Mokshetsky, Sigismond, 247–48.
Morrill, A. W., 268–73, 342–48.
Mosher, F. H., 368–73.
Mosquito (anti) convention of New Jersey, 244–46.
Muir, F., 459–60.
Musca domestica, 281–89.
Myzus fragæfolii, 135.
Myzocallis quercus, 130.

Naphthaline, 200.
Newell, Wilmon, 87–91, 92–96, 147, 153.
New Jersey anti-mosquito convention, 244–46.
Nezara hilaris, 336–39.
viridula, 336.
Nicine, 198.
Notolophus antiqua, 370, 371.
Nun moth, 474.

O'Kane, W. C., 7–22, 181–83.
Onion maggot, 197–200.
thrips, 196–97.

Orton, W. A., 109–16.
Orthezia insignis, 471.
Osborn, Herbert, 192.

Paddock, F. B., 183–88.
Palmer worm, 191–92.
Parker, J. R., 136–41.
Parks, T. H., 297, 417–21.
Parlatoria blanchardii, 243.
Parrott, P. J., 50–64, 227–28, 487.
Peach, arsenate injury, 165.
borer, lesser, 401–403.
Peairs, L. M., 152–53, 174–79.
Pear thrips, 478–79.
Pecan twig girdler, 218.
Pellegra, 293–94.
Pemphigus betæ, 136–41.
californicus, 127.
Perissopterus pulchellus, 227.
Perkinsiella saccharicida, 243.
Peruvian coccidæ, 467–73.
Phænicococcus marlatti, 243.
Pheidole megacephala, 396.
Phlebotomus verrucarum, 357.
Phorodon humuli, 440, 441, 443.
Phylloxera vastatrix, 297.
Phytonomus posticus, 417–21.
Picea englemanni, 246.
Pine moth, 340.
Pissodes strobi, 375.
Plant louse notes, 127–36.
quarantine, 109–16.
Plodia interpunctella, 204, 318.
Plum curculio, 168.
Poison resistance, 170.
Poisoned bran, 202.
Pokeweed, 327.
Pollenia rudis, 487.
Polygnotus hiemalis, 297.
minutus, 296, 297.
Popenoe, E. A., 155.
Porosagrotis delorata, 201–203.
orthogonia, 203.
Porthetria dispar, 178, 371.
Porto Rican sugar cane insects, 461–63.
Posthörner, 340.
Potassium sulphid, 335.
Potato weevil, 242.
Prenes ares, 462.
Prospaltella berlesei, 298.
perniciosi, 227, 422–32.
Pseudanthonomus validus, 193–94.
Pseudaonidia articulatus, 471.
Pseudococcus calceolariæ, 460.
citri, 148, 472.
sacchari, 461.
Pseudokermes cooleyi, 246–47.
Pulvinaria sp., 472.

Red spider, 324–36.
Reviews, 350, 408, 483.
Rhabdocnemis obscurus, 390, 444, 455.
Rhagoletis pomonella, 398–400.
Rhigopsidius tucumanus, 242.
Rhopalosiphum corylinum, 134.
Rhipiphorus pectinatus, 383, 387.
Ribes setosum, 194, 195.
Rice water weevil, 433–39.
Rice weevil, 204, 206–10, 315.
Riley, W. A., 157.
Rogers, D. M., 116–19.
Rohwer, S. A., 479–81.
Rose, agrilus infesting, 439.
Rotation, 333.
Rust, E. W., 467.

Saissetia hemisphærica, 471.
oleæ, 471.
San José scale, 141–47, 167–71, 422.
parasites, 227–28, 229.
Sand fly, 293–94.
Sanders, J. G., 102–106.
Sasscer, E. R., 240–44.
Scarabæid larvæ rearing, 445.
Scientific notes, 154, 250, 297, 339, 348, 405, 483.
Scolothrips sexmaculata, 331.
Scott, W. M., 478–79.
Serpentine leaf miner, 447–53.
Shelford, V. E., 229–33, 249.
Sherman, Franklin, 348.
Signiphora nigrita, 227.
Silvanus surinamensis, 204, 316.
Simulium vittatum, 293.
Sitotroga cerealella, 204.
Sitodrepa panicea, 317.
Slingerland, M. V., 408.
Smynthurus sp., 400–401.
Soap wash, 198.
Sodium cyanide, 307.
Soil fumigation, 305–12.
Spaulding, Perley, 106–108.
Spilogale interruptans, 153.
putorius, 252.
Sphenophorus nebulosus, 445–46.
Sporotrichum globuliferum, 474.
Spring-tails, 400–401.
Sterility in oats caused by thrips, 211–18.
Stethorus punctum, 330.
Stomoxys calcitrans, 459–60.
nebulosa, 459.
pallida, 459.
Strawberry, 479.
Sugar beet root louse, 136–41.
cane insects, 444, 461–62.
cane weevil borer, 455.
maples, 173.
Sulphur, 188.
Swezey, O., 455–57.
Synanthedon pictipes, 401.
Syrphus americanus, 295, 296.

Tachinid parasites, 390–98, 455.
Tæniothrips pyri, 478–79.
Tapinostola musculosa, 298.
Tarsonemus spinipes, 463.
Tartar, H. V., 463–67.

Temperature, high, 313–22.
and oviposition, 417.
relation to insect development, 174–79.
Tenebrio molitor, 177.
Tenebrioides mauritanicus, 318.
molitor, 318.
Tetranychus bimaculatus, 324, 481.
Tetrastichus asparagi, 296.
Thrips cerealium, 212.
tabaci, 196–97.
Tiphia inornata, 382–89.
Tolype laricis, 371.
Tower, D. G., 422–32.
Townsend, C. H. T., 357–67.
Toxoptera graminum, 403–404, 414, 415.
Tribolium confusum, 204, 316, 318.
ferrugineum, 204.
Triphleps insidiosus, 330.
Trogoderma ornatum, 317.
Trombidium sp., 152.

Vaccinium species, 398.
Verruga work, progress, 357–67.

Walking stick, 299.
Warble fly, occurrence in Canada, 195.
Washburn, F. L., 34, 64, 348.
and Spangler, A. J., 119–23.
Webb, J. L., 432–38.
Webber, R. T., 368–73.
Webster, F. M., 403–404.
Webster, R. L., 33.
Weiss, H. B., 250–51, 438–40.
Wheat stem maggot, 212.
White grubs, 382, 386, 389.
pine weevil, 375.
Whitmarsh, R. D., 336–39.
Wire worms, 308.
Wolcott, G. N., 382–89.
Woods, W. C., 398–400.

Xylina antennata, 189–91.

Ypsolophus ligulellus, 191–92.
pometellus, 191–92.

Zacher, F., 483.
Zagrammosoma multilineata, 454.
Zelus rubidus, 462.
Zenoleum, 286, 287.
Zoöphagous itonididæ, 458–59.